



# Emergent Phenomena in QCD & at the LHC

Peter Skands (Monash University)



Physics Colloquium, Melbourne University  
25 November 2014

# Emergence

The **elementary** interactions are encoded in the **Lagrangian**  
QFT → Feynman Diagrams → Perturbative Expansions (in  $\alpha_s$ )

The emergent is unlike its components insofar as ... it cannot be reduced to their sum or their difference."

*G. Lewes (1875)*

## ***Emergent phenomena in QCD***

Cannot be guessed directly from Lagrangian.

Two sources of emergence in QCD:

1. Scale Invariance (*can actually be guessed*)
2. Confinement (*win \$1,000,000 if you can prove*)


# The Constituents of QCD

The **elementary** interactions are encoded in the **Lagrangian**  
 QFT → Feynman Diagrams → Perturbative Expansions (in  $\alpha_s$ )

$$g_s^2 = 4\pi\alpha_s$$

THE BASIC ELEMENTS OF QCD: QUARKS AND GLUONS

$$\psi_q^j = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix}$$



$$\mathcal{L} = \bar{\psi}_q^i (i\gamma^\mu) (D_\mu)_{ij} \psi_q^j - m_q \bar{\psi}_q^i \psi_{qi} - \frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

$$D_{\mu ij} = \delta_{ij} \partial_\mu - ig_s T_{ij}^a A_\mu^a$$

Gauge Covariant Derivative: makes  $L$  invariant under  $SU(3)_C$  rotations of  $\psi_q$

$m_q$ : Quark Mass Terms  
(Higgs + QCD condensates)

Gluon-Field Kinetic Terms  
and Self-Interactions

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f^{abc} A_\mu^b A_\nu^c$$



There are more things in heaven and earth, Horatio, than are dreamt of in your philosophy

W. Shakespeare, Hamlet.

AUTUMN (in Australia)

LHC RUN 2: STARTS ~~SPRING~~ 2015!!!

ALMOST TWICE THE ENERGY (13 TeV compared with 8 TeV) AND MORE INTENSE BEAMS

$$\mathcal{L} = \bar{\psi}_q^i (i\gamma^\mu) (D_\mu)_{ij} \psi_q^j - m_q \bar{\psi}_q^i \psi_{qi} - \frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} \\ + \dots \dots \dots ?$$

LHC: still no explicit “new physics”

→ we’re still looking for *deviations* from SM




Accurate modeling of QCD improve searches & precision



# ATLAS EXPERIMENT

Run Number: 162620, Event Number: 16060241

Date: 2010-08-24 19:45:23 CEST



- 1st jet:  $p_T = 520$  GeV,  $\eta = -1.4$ ,  $\phi = -2.0$
- 2nd jet:  $p_T = 460$  GeV,  $\eta = 2.2$ ,  $\phi = 1.0$
- 3rd jet:  $p_T = 130$  GeV,  $\eta = -0.3$ ,  $\phi = 1.2$
- 4th jet:  $p_T = 50$  GeV,  $\eta = -1.0$ ,  $\phi = -2.9$

# QCD in the Ultraviolet

The “running” of  $\alpha_s$ :

$$Q^2 \frac{\partial \alpha_s}{\partial Q^2} = -\alpha_s^2 (b_0 + b_1 \alpha_s + b_2 \alpha_s^2 + \dots),$$

$$b_0 = \frac{11C_A - 2n_f}{12\pi} \quad C_A=3 \text{ for SU(3)}$$

$$b_1 = \frac{17C_A^2 - 5C_A n_f - 3C_F n_f}{24\pi^2} = \frac{153 - 19n_f}{24\pi^2}$$

$$b_2 = \frac{2857 - 5033n_f + 325n_f^2}{128\pi^3}$$

$b_3 = \text{known}$


At high scales  $Q \gg 1 \text{ GeV}$

Coupling  $\alpha_s(Q) \ll 1$

**Perturbation theory** in  $\alpha_s$  should be **reliable**: LO, NLO, NNLO, ...

E.g., in event shown on previous slide:

- 1st jet:  $p_T = 520 \text{ GeV}$
- 2nd jet:  $p_T = 460 \text{ GeV}$
- 3rd jet:  $p_T = 130 \text{ GeV}$
- 4th jet:  $p_T = 50 \text{ GeV}$



Full symbols are results based on N3LO QCD, open circles are based on NNLO, open triangles and squares on NLO QCD. The cross-filled square is based on lattice QCD.

# The Infrared Strikes Back

Naively, QCD radiation suppressed by  $\alpha_s \approx 0.1$

Truncate at fixed order = LO, NLO, ...

E.g.,  $\sigma(X+\text{jet})/\sigma(X) \propto \alpha_s$

**Example:** Pair production of SUSY particles at LHC<sub>14</sub>, with  $M_{\text{SUSY}} \approx 600$  GeV

LHC - sps1a - m~600 GeV

Plehn, Rainwater, PS PLB645(2007)217

FIXED ORDER pQCD	$\sigma_{\text{tot}}$ [pb]	$\tilde{g}\tilde{g}$	$\tilde{u}_L\tilde{g}$	$\tilde{u}_L\tilde{u}_L^*$	$\tilde{u}_L\tilde{u}_L$	$TT$
$p_{T,j} > 100$ GeV	$\sigma_{0j}$	4.83	5.65	0.286	0.502	1.30
inclusive X + 1 "jet"	$\sigma_{1j}$	2.89	2.74	0.136	0.145	0.73
inclusive X + 2 "jets"	$\sigma_{2j}$	1.09	0.85	0.049	0.039	0.26
$p_{T,j} > 50$ GeV	$\sigma_{0j}$	4.83	5.65	0.286	0.502	1.30
	$\sigma_{1j}$	5.90	5.37	0.283	0.285	1.50
	$\sigma_{2j}$	4.17	3.18	0.179	0.117	1.21

(Computed with SUSY-MadGraph)

$\sigma$  for X + jets much larger than naive estimate

$\sigma_{50} \sim \sigma_{\text{tot}}$  tells us that there will "always" be a ~ 50-GeV jet "inside" a 600-GeV process

All the scales are high,  $Q \gg 1$  GeV, so perturbation theory **should** be OK ...

# Conformal QCD



The Lagrangian of QCD is **scale invariant**

(neglecting small quark masses)

Characteristic of point-like constituents

To first approximation, observables depend only on dimensionless quantities, like **angles** and energy **ratios**

Also means that when we look closer at these constituents, they must generate ever self-similar patterns = fractals



James Bjorken  
"Lightcone Scaling"  
aka Bjorken Scaling;  
Conformal invariance

Note: scaling **violation** is induced in full QCD, but only by renormalization:  $g_s^2 = 4\pi\alpha_s(\mu)$

# (some) Physics

cf. equivalent-photon  
approximation  
Weiszäcker, Williams  
~ 1934

Charges Stopped  
or kicked

Radiation


Radiation

a.k.a.  
Bremsstrahlung  
Synchrotron Radiation

The harder they stop, the harder the  
fluctuations that continue to become radiation

# Jets $\approx$ Fractals

- Most bremsstrahlung is driven by divergent propagators  $\rightarrow$  simple structure
- Amplitudes factorize in singular limits ( $\rightarrow$  universal “conformal” or “fractal” structure)



Partons  $ab \rightarrow$   
“collinear”:

$P(z)$  = DGLAP splitting kernels, with  $z$  = energy fraction =  $E_a/(E_a+E_b)$

$$|\mathcal{M}_{F+1}(\dots, a, b, \dots)|^2 \xrightarrow{a||b} g_s^2 C \frac{P(z)}{2(p_a \cdot p_b)} |\mathcal{M}_F(\dots, a + b, \dots)|^2$$

Coherence  $\rightarrow$  Parton  $j$  really emitted by  $(i, k)$  “colour antenna” (in leading colour approximation)

Gluon  $j$

$\rightarrow$  “soft”:


$$|\mathcal{M}_{F+1}(\dots, i, j, k, \dots)|^2 \xrightarrow{j_g \rightarrow 0} g_s^2 C \frac{(p_i \cdot p_k)}{(p_i \cdot p_j)(p_j \cdot p_k)} |\mathcal{M}_F(\dots, i, k, \dots)|^2$$

+ scaling violation:  $g_s^2 \rightarrow 4\pi\alpha_s(Q^2)$

Can apply this many times  $\rightarrow$  nested factorizations  
**Jets-within-jets-within-jets ...**

# From Legs to Loops

**Unitarity:**  $\text{sum}(\text{probability}) = 1$



- $\rightarrow$  Can also include loops-within-loops-within-loops ...**
- $\rightarrow$  Bootstrap for approximate All-Orders Quantum Corrections!**

**Parton Showers:** reformulation of pQCD corrections as gain-loss diff eq.

Iterative (Markov-Chain) evolution algorithm, based on universality and unitarity

With evolution kernel  $\sim \frac{|\mathcal{M}_{n+1}|^2}{|\mathcal{M}_n|^2}$  (or soft/collinear approx thereof)

Generate explicit fractal structure across all scales (via Monte Carlo Simulation)

Evolve in some measure of *resolution*  $\sim$  hardness, virtuality,  $1/\text{time} \dots \sim$  fractal scale

+ account for scaling violation via quark masses and  $g_s^2 \rightarrow 4\pi\alpha_s(Q^2)$

# Our Research



## Parton Showers are based on $1 \rightarrow 2$ splittings


I.e., each **parton** undergoes a sequence of splittings

Multi-parton coherence effects can be included via “angular ordering”

Or via “dipole radiation functions”

(~ partitions dipole radiation pattern into 2 monopole terms)

Recoil effects needed to impose (E,p) cons (“local” or “global”)



E.g., **PYTHIA** (also HERWIG, SHERPA)

## At Monash, we develop an **Antenna Shower**, in which splittings are fundamentally $2 \rightarrow 3$

Each colour **dipole** undergoes a sequence of splittings

+ Intrinsically includes dipole coherence (leading  $N_c$ )

+ Lorentz invariance and explicit local (E,p) conservation


+ The non-perturbative limit of a colour dipole is a string piece

**Roots in Lund ~ mid-80ies: Gustafson & Petterson, Nucl.Phys. B306 (1988) 746**

What’s new in our approach?

Higher-order perturbative effects can be introduced via calculable corrections in an elegant and very efficient way

+ Writing a genuine antenna shower also for the initial state evolution



E.g., **VINCIA**  
(also ARIADNE)



Cf a lattice and its dual lattice  
Can either perceive of lattice **sites**  
or lattice **links**. Equivalent (dual) representations.

# VINCIA: Markovian pQCD\*

\*)pQCD : perturbative QCD

Start at Lowest Order

$$|M_F|^2$$

Generate "shower" emission

$$|M_{F+1}|^2 \stackrel{LL}{\sim} \sum_{i \in \text{ant}} a_i |M_F|^2$$

Correct to Matrix Element

$$a_i \rightarrow \frac{|M_{F+1}|^2}{\sum a_i |M_F|^2} a_i$$


Unitarity of Shower

$$\text{Virtual} = - \int \text{Real}$$

Correct to Matrix Element

$$|M_F|^2 \rightarrow |M_F|^2 + 2\text{Re}[M_F^1 M_F^0] + \int \text{Real}$$

Repeat



"Higher-Order Corrections To Timelike Jets"  
GeeKS: Giele, Kosower, Skands, PRD 84 (2011) 054003

# Quo Vadis?

## All sights are on **Run 2 of the LHC**

Next order of precision for jet rates and structure

Aid precision measurements and enhance discovery reach

Vast multi-jet phase spaces to explore with LHC

+ higher calculational efficiencies : SPEED

(has become a major issue for highly complicated final states)

Test runs in  $e^+e^-$  show factors  $10^2 - 10^3$  increases over conventional schemes

+ systematic and automated theory uncertainties

Part of being precise is knowing **how** precise. Our job to give an answer.



## Understanding the fractal

Unitarity and the structure of perturbative QCD

Beyond the Leading-Logarithmic approximation?

Beyond the Leading-Colour approximation?

The Structure of the proton (parton distributions)



# + Applications

## Example: The Top Quark

Heaviest known elementary particle:

$$m_t \sim 187 u (\sim m_{Au})$$

$$\text{Lifetime: } 10^{-24} \text{ s}$$

Complicated decay chains:

$$t \rightarrow bW^+ \quad \bar{t} \rightarrow \bar{b}W^-$$

$$W \rightarrow \{q\bar{q}', l\nu\}$$

quarks  $\rightarrow$  jets


b-quarks  $\rightarrow$  b-jets

$$m_t^2 \approx (p_b + p_{W^+})^2$$

$$\approx (p_{b\text{-jet}} + p_{q\text{-jet}} + p_{\bar{q}\text{-jet}})^2$$

**Accurate** jet energy calibrations  $\rightarrow m_t$

**Analogously** for any process / measurement involving coloured partons




# Long Wavelengths $> 10^{-15}$ m

## Quark-Antiquark Potential

As function of separation distance

What physical system has a linear potential?



$$F(r) \approx \text{const} = \kappa \approx 1 \text{ GeV/fm} \iff V(r) \approx \kappa r$$

~ Force required to lift a 16-ton truck

# String Breaks

In QCD, strings can (and do) break!

(In superconductors, would require magnetic monopoles)

In QCD, the roles of electric and magnetic are reversed

Quarks (and antiquarks) are “chromoelectric monopoles”

There are at least two possible analogies ~ tunneling:

CANONICAL

**Schwinger Effect**

Non-perturbative creation of  $e^+e^-$  pairs in a strong external Electric field

Probability from Tunneling Factor

$$\mathcal{P} \propto \exp\left(\frac{-m^2 - p_{\perp}^2}{\kappa/\pi}\right)$$

( $\kappa$  is the string tension equivalent)

**Hawking Radiation**

Non-perturbative creation of radiation quanta in a strong gravitational field

Thermal (Boltzmann) Factor



$$\mathcal{P} \propto \exp\left(\frac{-E}{k_B T_H}\right)$$

Linear Energy Exponent

ALTERNATIVE?


# The Lund String

- **Quarks** → String Endpoints
- **Gluons** → Transverse Excitations (kinks)



Gluon = kink on string, carrying energy and momentum

String Breaks by Tunneling (Schwinger Type)




- Probability of string break constant per unit area → **AREA LAW**
- Breakup vertices causally disconnected → order is irrelevant → iterative algorithm

# Colour Confusion

## Between which partons do confining potentials arise?

$e^+e^-$  : too easy



(still quite simple even after including bremsstrahlung etc.)

At  $e^+e^-$  colliders (eg LEP) - We generally find quite good agreement between **measured** particle spectra and **models** based on parton/antenna showers + strings

(with a couple of interesting exceptions, not covered here)

“Leading Colour” dipole decomposition works well


→ re-use same models as input for LHC (universality) ?

Proton-Proton (LHC)


More colour kicked around (& also colour in initial state)

Include “Beam Remnants”

Still might look relatively simple, to begin with



But no law against *several* parton-parton interactions




In fact, can easily be shown to happen frequently  
Included in all (modern) Monte Carlo models  
But how to make sense of the colour structure?


# Collective Effects?

A rough indicator of how much colour gets kicked around, should be the number of particles produced


So we study event properties as a function of " $N_{ch}$ " =  $N_{tracks}$



Independent Particle Production:  
→ averages stay the same



Correlations / Collective effects:  
→ averages depend on  $N_{ch}$



Plot shows the average transverse momentum versus  $N_{ch}$

# What are “Colour Reconnections”?

Simple example:  $e^+e^- \rightarrow W^+W^- \rightarrow q_1\bar{q}_2q_3\bar{q}_4$

Intensely studied at LEP2.

CR implied a non-perturbative uncertainty on the W mass measurement,  $\Delta M_W \sim 40 \text{ MeV}$

CR constrained to  $\sim 10\% \sim 1/NC^2$

Simple two-string system. What about pp?

## Several modelling attempts

Based on minimising the string action

String interactions (Khoze, Sjostrand)

Generalized Area Law (Rathsman et al.)




Colour Annealing (Skands, Wicke)

Gluon Move Model (Sjostrand et al.)

Based on SU(3) group multiplet weights

Dipole Swing (Lonnblad et al.)  $3 \otimes \bar{3} = 8 \oplus 1$

Generalized colour coherence (Christensen, Skands, in progress)



$$3 \otimes 3 = 6 \oplus \bar{3}.$$

$$8 \otimes 8 = 27 \oplus 10 \oplus \bar{10} \oplus 8 \oplus 8 \oplus 1$$

$$3 \otimes 8 = 15 \oplus 6 \oplus 3,$$

# Collective Effects?

There is now quite a lot of confusion in the field

Old-fashioned string models don't work that well at LHC

Eg need "CR" and don't reproduce low- $p_T$  identified-particle spectra

Quark-gluon plasma inspired models?

Using hydrodynamics (eg EPOS)

Statistical (Thermal) Distributions

Good fits ... even for ee ... but ... thermal???

**And how to reconcile with string picture?**

Colour-(re)connection / String Effects?

Subleading colour effects?

**Multi-parton coherence?**


**Colour accidents?**

**Soft-gluon exchanges?**

String-string interaction effects?

More colour charge: strings with higher tension?

Rescattering Effects (parton-parton or hadron-hadron)



# Summary

## Jets

Discovered at SPEAR (SLAC '72) and DORIS (DESY '73):  $E_{CM} \sim 5 \text{ GeV}$   
Collimated sprays of nuclear matter (hadrons).

Quasi-fractal structure of jets-within-jets & loops-within-loops

Simulated by parton-, dipole-, or antenna showers

Complementary to standard (LO, NLO, ...) perturbative matrix elements

**Showers are most precise for relatively soft/collinear radiation**

**Fixed-order calculations are most precise for relatively “hard” radiation**

**Much focus on how to combine the two consistently and efficiently: “matching”**

Unitarity is a key aspect of both approaches; sums & detailed balance.

## Strings enforce confinement



~ well understood in “sparse” environments ~ vacuum

Many indications that confinement is more complicated in pp

LHC Run 1 provided a treasure trove of data.

We are learning which questions to ask; what to measure in **Run 2** !

# Summary




QCD spans a huge variety of phenomena

Collective Effects

Amplitudes

Confinement


$$\mathcal{L} = \bar{\psi}_q^i (i\gamma^\mu) (D_\mu)_{ij} \psi_q^j - m_q \bar{\psi}_q^i \psi_{qi} - \frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$



Ultra-High Energies

Still only partially solved

Fragmentation



# Come to Australia



$P$



Establishing a new research direction in **Melbourne**  
Working on Precision LHC **phenomenology & soft physics**


**PYTHIA & VINCIA**

**NLO** Event Generators

Support LHC **experiments**, **astro-particle** community, and **future** accelerators

Outreach and Citizen Science

$P$



**Now Advertising** (on inSPIRE, AJO, ...):  
1 post doc in high-energy phenomenology  
2 PhD scholarships in theoretical physics  
(1 joint with Warwick ATLAS group, UK)  
+ Monash scholarships

**Oct 2014**  
→ Monash University  
Melbourne, Australia

# Asymptotic Freedom

“What this year's Laureates discovered was something that, at first sight, seemed completely contradictory. The interpretation of their mathematical result was that the closer the quarks are to each other, the *weaker* is the 'colour charge'. When the quarks are really close to each other, the ~~force~~<sup>charge</sup> is so weak that they behave almost as free particles. This phenomenon is called ‘asymptotic freedom’. The converse is true when the quarks move apart: the ~~force~~<sup>potential</sup> becomes stronger when the distance increases.”



The Nobel Prize in Physics 2004  
David J. Gross, H. David Politzer, Frank Wilczek



David J. Gross




H. David Politzer



Frank Wilczek

The Nobel Prize in Physics 2004 was awarded jointly to David J. Gross, H. David Politzer and Frank Wilczek "for the discovery of asymptotic freedom in the theory of the strong interaction".

Photos: Copyright © The Nobel Foundation



\*1 The force still goes to  $\infty$  as  $r \rightarrow 0$   
(Coulomb potential), just less slowly

\*2 The potential grows linearly as  $r \rightarrow \infty$ , so the force actually becomes constant  
(even this is only true in “quenched” QCD. In real QCD, the force eventually vanishes for  $r \gg 1 \text{ fm}$ )

# Evolution Equations

## What we need is a differential equation

Boundary condition: a few partons defined at a high scale ( $Q_F$ )

Then evolves (or “runs”) that parton system down to a low scale (the hadronization cutoff  $\sim 1$  GeV)  $\rightarrow$  It’s an evolution equation in  $Q_F$

## Close analogue: nuclear decay

Evolve an unstable nucleus. Check if it decays + follow chains of decays.

Decay constant

$$\frac{dP(t)}{dt} = c_N$$

Probability to remain undecayed in the time interval  $[t_1, t_2]$

$$\Delta(t_1, t_2) = \exp\left(-\int_{t_1}^{t_2} c_N dt\right) = \exp(-c_N \Delta t)$$

Decay probability per unit time

$$\frac{dP_{\text{res}}(t)}{dt} = \frac{-d\Delta}{dt} = c_N \Delta(t_1, t)$$

(requires that the nucleus did not already decay)


$$= 1 - c_N \Delta t + \mathcal{O}(c_N^2 \Delta t^2)$$

$\Delta(t_1, t_2)$  : “Sudakov Factor”

# Nuclear Decay

Nuclei remaining undecayed after time t

$$= \Delta(t_1, t_2) = \exp \left( - \int_{t_1}^{t_2} dt \frac{d\mathcal{P}}{dt} \right)$$



# The Sudakov Factor

In nuclear decay, the Sudakov factor counts:

How many nuclei remain undecayed after a time  $t$

Probability to remain undecayed in the time interval  $[t_1, t_2]$

$$\Delta(t_1, t_2) = \exp\left(-\int_{t_1}^{t_2} c_N dt\right) = \exp(-c_N \Delta t)$$

The Sudakov factor for a parton system counts:

The probability that the parton system doesn't evolve (branch) when we run the factorization scale ( $\sim 1/\text{time}$ ) from a high to a low scale

Evolution probability per unit "time"

$$\frac{dP_{\text{res}}(t)}{dt} = \frac{-d\Delta}{dt} = c_N \Delta(t_1, t) \quad \begin{array}{l} \text{(replace } t \text{ by shower evolution scale)} \\ \text{(replace } c_N \text{ by proper shower evolution kernels)} \end{array}$$

# What's the evolution kernel?

cf. conformal (fractal) QCD, Lecture 1  
(and PDF evolution, Lecture 2)


## DGLAP splitting functions

Can be derived from *collinear limit* of MEs  $(p_b+p_c)^2 \rightarrow 0$

+ evolution equation from invariance with respect to  $Q_F \rightarrow$  RGE

**DGLAP**  
(E.g., PYTHIA)

$$d\mathcal{P}_a = \sum_{b,c} \frac{\alpha_{abc}}{2\pi} P_{a \rightarrow bc}(z) dt dz .$$



$$p_b = z p_a$$

$$p_c = (1-z) p_a$$

$$P_{q \rightarrow qg}(z) = C_F \frac{1+z^2}{1-z} ,$$

$$P_{g \rightarrow gg}(z) = N_C \frac{(1-z(1-z))^2}{z(1-z)} ,$$

$$P_{g \rightarrow q\bar{q}}(z) = T_R (z^2 + (1-z)^2) ,$$

$$P_{q \rightarrow q\gamma}(z) = e_q^2 \frac{1+z^2}{1-z} ,$$

$$P_{l \rightarrow l\gamma}(z) = e_l^2 \frac{1+z^2}{1-z} ,$$

$$dt = \frac{dQ^2}{Q^2} = d \ln Q^2$$

... with  $Q^2$  some measure of "hardness"  
= event/jet resolution  
measuring parton virtualities / formation time / ...

**Note:** there exist now also alternatives to AP kernels (with same collinear limits!): dipoles, antennae, ...